


# Numerical model of a partially-ionized solar atmosphere

Kris Murawski,  
Group of Astrophysics, UMCS, Poland

September 25, 2017

- ▶ Our goals
- ▶ Two-fluid equations
- ▶ Two-fluid waves
- ▶ JOANNA code
- ▶ Case study - granulation
- ▶ Conclusions



## Our goals:

- ▶ Contribute to solving some problems for weakly ionized plasma (lower solar atmospheric layers, ionospheres/thermospheres of planets)
- ▶ Develop our own 2-fluid code

# Two-fluid equations

## Equations for neutrals

*Euler equations:*

$$\frac{\partial \rho_n}{\partial t} + \nabla \cdot (\rho_n \mathbf{V}_n) = -S_1, \quad (1)$$

$$\frac{\partial (\rho_n \mathbf{V}_n)}{\partial t} + \nabla \cdot (\rho_n \mathbf{V}_n \mathbf{V}_n) + \nabla p_n - \rho_n \mathbf{g} = -S_2, \quad (2)$$

$$\frac{\partial E_n}{\partial t} + \nabla \cdot ((E_n + p_n) \mathbf{V}_n) - \rho_n \mathbf{g} \cdot \mathbf{V}_n - q_n = -S_3. \quad (3)$$

# Two-fluid equations

## Equations for ions

*MHD equations:*

$$\frac{\partial \rho_i}{\partial t} + \nabla \cdot (\rho_i \mathbf{V}_i) = S_1, \quad (4)$$

$$\frac{\partial(\rho_i \mathbf{V}_i)}{\partial t} + \nabla \cdot (\rho_i \mathbf{V}_i \mathbf{V}_i) + \nabla p_i - \rho_i \mathbf{g} - \frac{1}{\mu} (\nabla \times \mathbf{B}) \times \mathbf{B} = S_2, \quad (5)$$

$$\frac{\partial E_i}{\partial t} + \nabla \cdot ((E_i + p_i) \mathbf{V}_i) - \rho_i \mathbf{g} \cdot \mathbf{V}_i - q_i = S_3, \quad (6)$$

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{V}_i \times \mathbf{B}) + \mathbf{S}_e, \quad \nabla \cdot \mathbf{B} = 0. \quad (7)$$

# Two-fluid equations

## Source terms

$$S_1 = -\rho_i(\alpha_r \rho_i - a_i \rho_n), \quad (8)$$

$$S_2 = a_c \rho_i \rho_n (\mathbf{V}_n - \mathbf{V}_i) - \rho_i (\alpha_r \rho_i \mathbf{V}_i - a_i \rho_n \mathbf{V}_n), \quad (9)$$

$$S_3 = a_c \rho_i \rho_n (\mathbf{V}_n - \mathbf{V}_i) \cdot \mathbf{V}_i. \quad (10)$$

$S_1$  - ionization/recombination,

$S_2$  - ion-neutral collisions,

$S_3$  - energy source term

(Smith & Sakai 2008, Zaqrashvili et al. 2011, 2012, Meier & Shumlak 2012).

# Two-fluid waves

- ▶ **HD waves:** 1 acoustic and 1 entropy mode (Goedbloed & Poedts 2004, Murawski et al. 2011)
- ▶ **MHD waves:** 1 Alfvén, 2 (slow and fast) magnetoacoustic, and 1 entropy mode
- ▶ **Two-fluid waves:**
  - MHD waves + extra entropy waves (Zaqarashvili et al. 2011, Soler et al. 2016)
  - ▶ dispersive, damped, for real  $\mathbf{k}$  cut-off for slow neutral wave
  - ▶ Effective damping of Alfvén and kink waves  $\rightarrow$  plasma heating

# Few remarks on waves

- ▶ Ion-neutral collisions introduce characteristic scales - waves become dispersive in a homogeneous medium
- ▶ Gravity introduces dispersion and cut-off (Lamb/Klein-Gordon equation)
- ▶ Shock - abrupt changes in all fluid quantities
- ▶ Pseudo-shock/entropy mode - sudden change in mass density alone, while other fluid quantities are smooth across this wave
- ▶ Rarefaction wave (nonlinear)




- ▶ Developed by Darek Wójcik
- ▶ Targets: HD, MHD, **2-fluid**, any-system of hyperbolic/parabolic eqs
- ▶ Multi-physics: non-adiabatic, non-ideal terms
- ▶ Shock-capturing algorithms: HLLC, HLLD, MUSTA
- ▶  $\nabla \cdot \mathbf{B}$  cleaning by GLM (Dedner et al. 2002)
- ▶ Reconstruction: flat, linear, PPM, WENO3
- ▶ More on <http://kft.umcs.lublin.pl/dwojcik/>.

# The hydrostatic solar atmosphere

Static equilibrium ( $\mathbf{V}_{i,n} = 0$ ) with force-free (current-free)  $\mathbf{B}$ ,

$$-\nabla p_{i,n} + \frac{1}{\mu}(\nabla \times \mathbf{B}) \times \mathbf{B} + \varrho_{i,n} \mathbf{g} = 0. \quad (11)$$



**Figure:** Solar temperature for ions (dashed line) and neutrals (dotted line) (Avrett & Loeser 2008, Wójcik 2017).

# Case study - solar granulation

Show the movie (courtesy of Darek Wójcik 2017).

# Conclusions

- ▶ Well tested JOANNA code - passed many (HD and MHD) tests (<http://kft.umcs.lublin.pl/dwojcik/>)
- ▶ Robust code - simulation of 2-fluid convection
- ▶ Versatile code - can be adopted to any hyperbolic/parabolic set of equations